

DELAYED RECOMBINATION

P. J. E. PEEBLES¹, S. SEAGER², AND WAYNE HU²

¹Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544

²Institute for Advanced Study, Olden Lane, Princeton, NJ 08540

Draft version February 1, 2008

ABSTRACT

Under the standard model for recombination of the primeval plasma, and the cold dark matter model for structure formation, recent measurements of the first peak in the angular power spectrum of the cosmic microwave background temperature indicate the spatial geometry of the universe is nearly flat. If sources of Ly α resonance radiation, such as stars or active galactic nuclei, were present at $z \sim 1000$ they would delay recombination, shifting the first peak to larger angular scales, and producing a positive bias in this measure of space curvature. It can be distinguished from space curvature by its suppression of the secondary peaks in the spectrum.

Subject headings: cosmology: cosmic microwave background

1. INTRODUCTION

The measurements of the anisotropies of the cosmic microwave background (the CMB) offer extraordinarily powerful tests of the relativistic Friedmann-Lemaître cosmological model and the nature of the early stages of cosmic structure formation (e.g. Jungman et al. 1996). Indeed, the recent detection of the first peak in the angular power spectrum of the CMB temperature indicates space is close to flat (Miller et al. 1999; Melchiorri et al. 1999; de Bernardis et al. 2000). The interpretation is quite indirect, however so we must seek diagnostics for possible complications.

There are relatively few physical effects that can *increase* the angular scale of the first peak. Because of this fact, the observed large angular scale of the peak is believed to strongly disfavour open universes. Of the fundamental cosmological parameters, only a Hubble constant well in excess of observations can substantially increase the scale of the peak in an open or flat universe (Hu & Sugiyama 1995).

One possibility is that some process at redshift $z \sim 1000$ delayed recombination of the primeval plasma. This would increase the sound horizon at last scattering and decrease the angular size distance to last scattering, moving the first peak of the CMB temperature fluctuation spectrum to smaller angular wavenumber (Hu & White 1996; Weller et al. 1999), and biasing the measure of space curvature to a too large (more positive) apparent value. Another consequence of delayed recombination is that it would suppress the secondary peaks due to an increase in the time available for acoustic oscillations to dissipate (Silk 1968; Hu & White 1996).

Because the first peak is not observed to have suffered substantial dissipation, recombination in the delayed model must be rapid compared with the cosmological expansion rate. Thus if recombination were delayed by ionizing radiation from decaying dark matter (e.g. Sarkar & Cooper 1983; Sciama et al. 1991; Ellis et al. 1992) or evaporating primeval black holes (Naselskij & Polonarëv 1987), or by thermal energy input from cosmic string wakes (Weller et al. 1999), the source would have to terminate quite abruptly and well before $z \sim 100$ when the universe starts to become optically thin even if the baryons

are fully ionized.

Here we consider a picture that more naturally allows rapid recombination: sources of radiation at $z \sim 1000$ that produce many more photons in the resonance Ly α line than ionizing photons, in the manner of a quasar. The Ly α photons would increase the population in the principal quantum number $n = 2$ levels of atomic hydrogen, increasing the rate of photoionization from $n = 2$ by the CMB. Since the rate of thermal photoionization from $n = 2$ varies rapidly with redshift at $z \lesssim 1000$, the delayed recombination is rapid, and the residual ionization can be small enough that the optical depth for Thomson scattering after recombination is well below unity. Thus the height of the first peak in the spectrum of CMB temperature fluctuations is little affected by the delayed recombination. The shift in the angular wavenumber at the first peak is modest also, even if early sources produce many Ly α photons per baryon, but the shift can be considerably larger than the projected precision of the measurements. Thus it is fortunate that we seem to have an unambiguous diagnostic for delayed recombination in the suppression of the secondary peaks.

After this work was substantially complete we learned that the recent measurements by the BOOMERanG experiment in fact favor a substantial suppression of the second peak (de Bernardis et al. 2000). We emphasize that there are many other ways to account for this effect, and that there is no evidence for the early source of Ly α radiation assumed in our picture. Within the conventional adiabatic CDM model the recombination history is well understood (Seager et al. 2000 and earlier references therein), so there is good reason to expect the residual fluctuations in the CMB may be related to the cosmology in a simple and computable way. But it is good science to bear in mind the possibility that Nature is more complicated than our ideas, a rule that has particular force in cosmology because of the limited empirical basis.

2. THE MODEL

Since the early source of radiation is purely conjectural, a simple model is appropriate. We assume the rate of production of Ly α resonance photons per unit volume (in

excess of those produced by the primeval plasma) is

$$dn_\alpha/dt = \epsilon_\alpha n_H H(t), \quad (1)$$

where n_H is the number density of hydrogen nuclei, $H(t) = \dot{a}/a$ is the expansion rate, and ϵ_α is a free parameter. The computation of the ionization history by Seager et al. (2000) is readily adjusted to take account of this new source term if it is approximately homogeneous. The results in Figure 1 assume the Hubble parameter is $h = 0.7$ in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$, the baryon density parameter is $\Omega_b h^2 = 0.02$ (Tytler et al. 2000) and the total matter density parameter is $\Omega_m = 0.25$. The ionization history is quite similar at $\Omega_m = 0.4$.

The optical depth for Thomson scattering subsequent to redshift z is

$$\tau = 63x \left(\frac{1 - Y_p}{0.76} \right) \left(\frac{\Omega_b h^2}{0.02} \right) \left(\frac{\Omega_m h^2}{0.122} \right)^{-1/2} \left(\frac{z}{10^3} \right)^{3/2}, \quad (2)$$

if the fractional ionization is constant at x at redshift less than z . Figure 1 shows that the delayed recombination through the redshift z_* where τ passes through unity is about as fast relative to the expansion rate as in the standard recombination model. Furthermore the decrease in z_* itself is approximately

$$\frac{z_*(\epsilon_\alpha)}{z_*(0)} = (1 + 3\epsilon_\alpha)^{-0.042}. \quad (3)$$

Sources of Ly α photons may also produce ionizing radiation. Since the rate of recombination of fully ionized baryons at the cosmic mean density is faster than the rate of expansion the ionization, x may be approximated by the equilibrium equation

$$\alpha n_H x^2 \simeq \epsilon_i H. \quad (4)$$

The definition of the parameter ϵ_i follows equation (1), and α is the recombination coefficient for principal quantum numbers $n \geq 2$. At this ionization the optical depth for Thomson scattering is

$$\tau \sim \epsilon_i^{1/2} \left(\frac{1 - Y_p}{0.76} \right)^{1/2} \left(\frac{\Omega_b h^2}{0.02} \right)^{1/2} \left(\frac{\Omega_m h^2}{0.122} \right)^{-1/4} \left(\frac{z}{10^3} \right)^{1.2}. \quad (5)$$




FIG. 1.— The ionization fraction, x , as a function of redshift for $\epsilon_i = 0$ and various values of ϵ_α .

Figure 2 shows in more detail the effect of the production of ionizing photons on the ionization history. If $\epsilon_i \gtrsim 1$ the slow decrease in the optical depth through $\tau \sim 1$ would remove the first peak in the anisotropy power spectrum, an effect the measurements indicate is unacceptable.

Sources of Ly α photons may also add energy to the CMB by inverse Compton scattering, but if the energy added is comparable to the energy in Ly α photons the effect on the y -parameter is small.

3. TEMPERATURE ANISOTROPY POWER SPECTRUM

In the adiabatic CDM model the ionization history and the cosmological parameters fix the CMB anisotropy power spectrum; the results in Figure 2 for the cosmologically flat model with the parameters in Figure 1, and in Figure 3 for an open model with the cosmological parameters $\Omega_m = 0.6$, $\Omega_\Lambda = 0$, $h = 0.7$, $\Omega_b h^2 = 0.02$ are computed using a code based on White & Scott (1996).

The shift in the angular wavenumber l_1 at the peak is noteworthy because l_1 is used to infer the space curvature. A general expression for its value in adiabatic models is (Hu & Sugiyama 1995, with $l_1 \approx 0.73\ell_p$)

$$\ell_1 \approx \frac{125}{\sqrt{\Omega_m + \Omega_\Lambda}} [1 + \ln(1 - \Omega_\Lambda)^{0.085}] \left(\frac{z_*}{10^3} \right)^{1/2}, \quad (6)$$

$$\times \left(\frac{1}{\sqrt{R_*}} \ln \frac{\sqrt{1 + R_*} + \sqrt{R_* + r_* R_*}}{1 + \sqrt{r_* R_*}} \right)^{-1},$$

where

$$\begin{aligned} r_* &= 0.042 m^{-1} (z_*/10^3), \\ R_* &= 30b(z_*/10^3)^{-1}, \\ z_*(0) &\approx 1008(1 + 0.00124b^{-0.74})(1 + c_1 m^{c_2}), \\ c_1 &= 0.0783b^{-0.24}(1 + 39.5b^{0.76})^{-1}, \\ c_2 &= 0.56(1 + 21.1b^{1.8})^{-1}, \end{aligned} \quad (7)$$

with $b \equiv \Omega_b h^2$ and $m \equiv \Omega_m h^2$. Although in isocurvature models ℓ_1 is larger by 50% or more, it scales with cosmological parameters in the same way.

The leading order dependence of the peak scale is then

$$\ell_1 \propto (\Omega_m + \Omega_\Lambda)^{-1/2} z_*^{1/2}, \quad (8)$$




FIG. 2.— The effect of new sources of ionizing photons on the recombination history. For each value of ϵ_i the upper curve is for $\epsilon_\alpha = 1$ and the lower curve for $\epsilon_\alpha = 0$.

such a 10% change in z_* shifts ℓ_1 by 5% and can compensate a 10% change in the spatial curvature $\Omega_m + \Omega_\Lambda$. In our model for delayed recombination with z_* given by eqn. (3), $\epsilon_\alpha = 1000$ can make an open universe with $\Omega_m = 0.6$ appear flat (see Figure 4).

Even in the context of cosmologically flat models, one might want to consider the possibility of delayed recombination. The observed peak at $\ell_1 = 197 \pm 12$ (95% CL, de Bernardis et al. 2000) favors either a slightly closed low density universe or one of high density ($\Omega_m h^2$). While flat models are certainly still consistent with the data, the expected increase in precision of the measurements could force one into accepting either a closed geometry, large Hubble constant or delayed recombination.

The amplitudes of the secondary peaks break this approximate degeneracy of delayed recombination with space curvature or the model for structure formation. While the promptness of the delayed recombination nearly preserves the first peak in the spectrum the secondary peaks are strongly suppressed because of the extra time the photons are allowed to diffuse (Silk 1968). As discussed in Hu & White (1996), the ratio of the first peak location to the location of the damping tail is a sensitive test of delayed recombination. An equally important but more subtle effect is that a delay in recombination raises the baryon-photon momentum density ratio R_* due to the redshifting of the photons. This effect suppresses the second peak and raises the third. Though an increase in the baryon density parameter also has this effect, it simultaneously reduces rather than increases the damping and can in principle be distinguished through the higher peaks.

Under standard recombination, the the ratio of power at the second versus first peak scales as

$$A_2(0) \equiv \frac{C_{\ell_2}}{C_{\ell_1}} \approx 0.7 \left[1 + \left(\frac{\Omega_b h^2}{0.016} \right)^4 \right]^{-1/4} 2.4^{n-1} \quad (9)$$

In our model, $A_2(\epsilon_\alpha) \approx A_2(0)z_*(\epsilon_\alpha)/z_*(0)$ such that delayed recombination is twice as effective in suppressing power at the second peak as it is at shifting the first peak. Observations currently favor $A_2 \sim 1/3$ (see Fig. 4) com-




FIG. 3.— The temperature anisotropy power spectrum ($\Delta T = [\ell(\ell+1)C_\ell/2\pi]^{1/2}$) for various values of the Ly α production parameter ϵ_α .

pared with ~ 0.5 in our fiducial Λ CDM model with standard recombination. Of course the tilt n and the baryon density can also lower this ratio.

Because of its cumulative effect over the secondary peaks, ϵ_α has statistically significant effects as long as it is greater than

$$\epsilon_{\min} = \left[\sum_{\ell=2}^{\ell_{\max}} (\ell + 1/2) \left(\frac{\partial \ln C_\ell}{\partial \epsilon} \right)^2 \right]^{-1/2}, \quad (10)$$

where $\ell_{\max}$ is the largest ℓ for which the measurements are cosmic variance limited. For $\ell_{\max} = 500$, $\epsilon_{\min} = 0.006$; for $\ell_{\max} = 1000$, $\epsilon_{\min} = 0.002$.

4. DISCUSSION

Our model for delayed recombination with $\epsilon_\alpha = 1$ yields a 5% shift of the first peak and a reduction of the secondary peak by 10%, an observationally interesting effect. The sources must have $\epsilon_i \ll 1$ (see Figure 2). This condition requires that the Ly α sources (perhaps hot stars or quasars) be surrounded by envelopes of neutral primeval material which strongly absorbs the ionizing radiation.

Our picture does not follow from the conventional CDM model for structure formation. Apparent problems with excess small-scale clustering in this model (Moore et al. 1999; Klypin et al. 1999) have motivated discussions of modifications (Spergel & Steinhardt 1999; Kamionkowski & Liddle 1999; Peebles 2000; Hu, Barkana, & Gruzinov 2000). These modifications would tend to delay the appearance of the first generation of gravitationally bound systems, however, which is in the opposite direction to what is postulated here. These lines of thought do not rule out early sources of Ly α photons, of course, but they do suggest one might best look for a source outside the adiabatic CDM model. Perhaps cosmic strings produced wakes that were subdominant to the primeval CDM density fluctuations in determining the mass fluctuation power spectrum but did produce occasional non-gaussian density fluctuations (Contaldi et al. 1999) large enough to have




FIG. 4.— A delay in recombination (with $\epsilon_\alpha = 1000$) can make an open universe ($\Omega_m = 0.6$, dashed line) appear flat (solid line, $\Omega_m = 0.25$, $\Omega_\Lambda = 0.75$) or decrease ℓ_1 in a flat universe and suppress the secondary peaks (with $\epsilon_\alpha = 10$). The COBE and BOOMERanG data are plotted as bandwidth $\times 1\sigma$ error boxes.

collapsed to stars or active black holes that could have produced Ly α photons.

To summarize, we have argued that the picture of delayed recombination caused by early sources of Ly α radiation, modeled after quasar spectra, but with suppressed X-ray emission, has the virtue that it naturally preserves the observed first peak in the CMB temperature angular power spectrum while shifting the peak to larger scales. The picture is ad hoc but important as a conceivable complication in the application of an exceedingly powerful cosmological test. The most direct diagnostic for delayed recombination seems to be the suppression of the secondary peaks in the spectrum. The amplitude of the third peak, which may be measured by the BOOMERanG experiment or the MAP satellite¹, is crucial for distinguishing the effect of delayed recombination from that of adjustments of the values of space curvature, the baryon density, the Hubble constant or the shape of the spectrum of the primeval density fluctuations. If the measured secondary peaks agreed with the standard model for recombination with astronomically acceptable cosmological parameters that fit the primary peak, it would convincingly rule out our model for delayed recombination.

We have benefitted from discussions with Martin Rees. P.J.E.P. is supported in part by the NSF. S.S. is supported by NSF grant PHY-9513835. W.H. is supported by the Keck Foundation and a Sloan Fellowship.

REFERENCES

- Contaldi, C., Hindmarsh, M., & Magueijo, J. 1999, *Phys. Rev. Lett.*, 82, 2034
 de Bernardis, P. et al., 2000, *Nature*, 404, 955
 Ellis, J. et al. 1992, *Nuc. Phys. B*, 373, 399
 Hu, W., Barkana, R., & Gruzinov, A. 2000, preprint, astro-ph/0003365
 Hu, W. & Sugiyama N. 1995, *PRD* 51, 2599
 Hu, W. & White, M. 1996, *ApJ*, 471, 30
 Jungman, G., Kamionkowski, M., Kosowsky, A., Spergel, D. N., 1996, *Phys. Rev. D*, 54, 1332
 Kamionkowski, M. & Liddle, A.R. 1999, *Phys. Rev. Lett.*, submitted (astro-ph/9911103)
 Klypin, A., Kravtsov, A.V., Valenzuela, O. & Prada, F. 1999, *ApJ*, 522, 82
 Melchiorri, A. et al. 1999, preprint, astro-ph/9911445
 Miller, A.D. et al. 1999, *ApJ*, 524, 1
 Moore, B. et al. 1999, *ApJ*, 524, L19
 Nasel'skij, P. D. & Polonarov, A. G. 1987, *Soviet Astron Lett* 13, 67
 Peebles, P.J.E. 2000, preprint, astro-ph/0002495
 Sarkar, S. & Cooper, A. M. 1983, *Phys. Lett. B*, 148, 347
 Seager, S., Sasselov, D. D. & Scott, D. 2000, *ApJ*, in press (astro-ph/9912182)
 Scott, D., Rees, M. J., & Sciama, D. W. 1991, *AA*, 250, 295
 Spergel, D. N. & Steinhardt, P. J. 1999, *Phys. Rev. Lett.*, in press (astro-ph/9909386)
 Tytler, D., O'Meara, L. M., Suzuki, N., & Lubin, D. 2000, *Physica Scripta*, astro-ph/0001318
 Weller, J., Battye, R. A., & Albrecht, A. 1999, *Phys. Rev. D* 60, 103520
 Silk, J. 1968, *ApJ* 151, 459
 White, M. & Scott, D. 1996, *ApJ*, 459, 415

¹<http://map.gsfc.nasa.gov>